TOBACCO USE

Healthy People 2020 Leading Health Indicators (LHIs)

Goal: Reduce illness, disability, and death related to tobacco use and secondhand smoke exposure.

TU-1.1 Reduce cigarette smoking by adults

BASELINE 2008 **20.6%**

2015 **15.1%**

of **adults** aged 18 years and older were current cigarette smokes in 2015

(age adjusted to the year 2000 standard population)

Data Source: National Health Interview Survey

TARGET **12%** YEAR 2020

TU 2.2 Reduce use of cigarettes by adolescents (past month)

BASELINE 2008 **19.5%**

2015 **10.8%**

of **adolescents** in grades 9 through 12 smoked cigarettes in the past 30 days in 2015

Data Source: Youth Risk Behavior Surveillance System

TARGET **16%** YEAR 2020

TU-11.1 Reduce the proportion of children aged 3 to 11 years exposed to secondhand smoke

BASELINE 2005-2008 **52.2%**

2009-2012 **41.3%**

of **children** aged 3 to 11 years were exposed to secondhand smoke in 2009 - 2012

Data Source: National Health and Nutrition Examination Survey

TARGET **47%** YEAR 2020

For more information, please visit **http://www.cdc.gov/tobacco/hp2020**

Email questions to: hp2020tobacco@cdc.gov

USE YOUR **SMARTPHONE** TO SCAN THIS **QR CODE**

Centers for Disease Control and Prevention
National Center for Chronic Disease Prevention and Health Promotion

CS270709-A OCTOBER 2016